

SERPENTINE : A MODEL HABITAT FOR BOTANICAL RESEARCH IN SRI LANKA.

Nishanta Rajakaruna¹ and Alan J. M. Baker²

¹Department of Biological Sciences, Stanford University, Stanford, CA 94305-5020, USA.

²School of Botany, The University of Melbourne, Parkville, VIC 3010, Australia

ABSTRACT

Serpentine (ultramafic) outcrops are found in most parts of the world and for many years have attracted the attention of soil scientists and plant biologists alike due to the unusual nature of both the substrate and the life forms they harbour. In this paper, we introduce serpentine outcrops as a model habitat for botanical studies and review preliminary findings of a recently conducted soil and floristic survey of four of the five known serpentine outcrops in Sri Lanka. We hope that this paper will stimulate much-needed studies, both exploratory and experimental, and point to the immediate need for conservation of this unique habitat.

INTRODUCTION

The study of plants growing under unusual soil conditions has a long and fascinating history (Kruckeberg, 2002). Early botanists and explorers often observed unique soil-plant associations and such 'soil-indicator plants' became useful in prospecting for natural deposits of valuable metals and minerals (Brooks, 1972; Baker and Brooks, 1988; Brooks and Johannes, 1990; Brooks *et al.*, 1995; Cannon, 1960, 1971). In the last decades there has been a growing interest in the use of edaphically restricted plants for the study of plant physiology, ecology, and evolution (Mason, 1946; Antonovics *et al.*, 1971; Proctor and Woodell, 1975; Brooks, 1987; Baker *et al.*, 1992; Macnair and Gardner, 1998; Balkwill, 2001). More recently, academic and industrial scientists have come together to explore how such plants can be used for both the reclamation of vast areas of contaminated soils (*i.e.*, phytoremediation) and the extraction of toxic metals (*i.e.*, phytomining) from metal-rich sites such as abandoned mine tailings and waste sites (Brooks, 1998; Brooks *et al.*, 1998; Whiting *et al.*, 2002, 2004; Reeves, 2003). All-in-all, the study of plants growing under extreme soil conditions holds much promise, providing both a model for botanical studies as well a means for an alternative to environmental rehabilitation.

In this regard, plants that grow on serpentine soils have attracted much attention from botanists in many parts of the world (Brooks, 1987; Baker *et al.*, 1992; Roberts and

Proctor, 1992; Balkwill, 2001). Serpentine soils are derived from serpentinite and other such ultramafic rocks and are located along continental margins and on offshore islands, often associated with obducted oceanic crust (ophiolite suites) and plate tectonics (Coleman and Jove, 1992). Ultramafic rocks occupy important positions in the geology of both Cuba and New Caledonia covering areas of 5000 km² and 5500 km², respectively. In New Caledonia, approximately one third of the landmass is occupied by rocks of ultramafic origin (Brooks, 1987). The Great Dyke of Zimbabwe is an extensive formation of serpentinite and related rocks about 500 km long and averaging 8 km wide (Wild and Bradshaw, 1977). In western North America, both local and massive occurrences of ultramafics are found; 2860 km² in California, 1170 km² in Oregon, and 520 km² in Washington (Kruckeberg, 1984). There are equally large areas of ultramafics in central Brazil (Brooks *et al.*, 1990). An extensive ultramafic rock cover of 8000 km² is found in Sulawesi, Indonesia (Baker *et al.*, 1992), however, the ultramafic regions of Asia are generally the least explored and poorly described of such regions in the world (Baker and Brooks, 1988; Proctor, 2003). Until recently, it was not known whether there were any occurrences of ultramafic rocks in Sri Lanka. Geological exploration work conducted in the 1970s revealed rocks of this type (Dissanayake and Van Riel, 1978; Munasinghe and Dissanayake, 1979, 1980; Dissanayake, 1982) located along a Precambrian suture zone between two plate boundaries, the

Vijayan and Highland Series. Since the deposits were northerly spread close to the boundary of these two Series and were fault-associated, it was believed that the serpentinites of Sri Lanka have a deep-seated origin. The work reported on the composition of the rocks suggests that the mineral sequences favour a primarily pyroxene-derived serpentinite (Dissanayake and Van Riel, 1978; Ranasinghe, 1987).

The nature of serpentine soils

Soils that weather from serpentinite and other ultramafic rocks are generally shallow and rocky. They have extreme physical and chemical properties that strongly reflect the elemental composition of the parental rock. Iron (Fe), magnesium (Mg), and silicon (Si) are chief elements of the ferromagnesian minerals found in serpentine soils, with nickel (Ni), cobalt (Co), chromium (Cr) and manganese (Mn) often occurring in exceptionally high concentrations. Serpentine soils have high values of exchangeable Mg and exceptionally low values of exchangeable calcium (Ca) and are usually deficient in nutrients such as nitrogen (N), phosphorus (P), potassium (K), boron (B), and molybdenum (Mo). Cation exchange capacities (CEC) range from 5.2 – 43 Cmol⁽⁺⁾ / kg dry soil; pH values are often high and can range from 6.1 – 8.8 (Brooks, 1987; Kruckeberg, 1992). The low nutrient status, cation imbalances and high metal concentrations, along with high temperature effects, moisture stress, soil instability, and resulting biotic conditions, limit plant growth and survival in serpentine habitats (Tadros, 1957; Proctor and Woodell, 1975; Arianoutsou *et al.*, 1993).

The stressed and highly selective nature of serpentine habitats is undoubtedly a consequence of the interplay of various physical, chemical, and biotic factors. The combination of these factors, with their particular intensities, forms the feedback loop that is known as the 'serpentine syndrome' (Jenny, 1980). The causes of the 'serpentine syndrome' have been traced to imbalance of Ca and Mg (Vlamiš and Jenny, 1948; Walker, 1954; Walker *et al.*, 1955; Madhok and Walker, 1969; Proctor and Woodell, 1975; Nagy and Proctor, 1997), Mg toxicity (Novak, 1928; Proctor, 1970), heavy metal toxicity (Antonovics *et al.*, 1971; Proctor and McGowan, 1976; Brooks, 1987), or low levels of essential nutrients such as N and P (Gordon and Lipman, 1926; Proctor and Woodell, 1975; Brooks, 1987; Proctor, 2003). A vast amount of research has

been conducted world-wide on various aspects of plant life on ultramafic substrates (Whittaker *et al.*, 1954; Krause, 1958; Proctor and Woodell, 1975; Kruckeberg, 1984, 1992; Brooks, 1987; Roberts and Proctor, 1992; Balkwill, 2001; Proctor, 2003) allowing us to make some informed generalizations on plants found in these 'harsh' environments.

The nature of serpentine plants

Plants growing on serpentine soils are often morphologically and physiologically adapted to deal with the physical and chemical characteristics of this extreme edaphic environment. They are frequently dwarf and xeromorphic, with chlorotic, narrow, and glaucous leaves. They show strong sclerenchymatic development and possess enlarged root systems (Kruckeberg, 1984). These observations imply that the plants are perhaps well-equipped to deal with both water stress and nutrient stresses prevalent in these habitats. In fact, an ongoing comparative study of serpentine shrubs and their closest relatives growing off of the serpentine substrate clearly show that there are multiple traits relating to both functional morphology and physiology that differ between such species pairs, sometimes even between populations of the same species (Rajakaruna and Ackerly, unpublished data). An important aspect of plants growing in serpentine soils is their capacity to tolerate and even hyperaccumulate (Brooks, 1998; Reeves, 2003) toxic levels of heavy metals, notably nickel. Plants found in these environments have both external and internal mechanisms to tolerate high metal concentrations (Baker, 1987; Baker and Walker, 1990; Verkleij and Schat, 1990; Pollard *et al.*, 2002; Baker *et al.*, 2000). By avoiding excessive uptake of metal ions, adopting excretory mechanisms, and developing storage devices, metal-tolerant plants are able to survive in soils toxic to most vegetation. Studies have shown that mechanisms of metal tolerance may be different in different species and for different metals (Shaw, 1990; Tilstone *et al.*, 1997; Llugany *et al.*, 2003). The evidence to date suggests that cell-wall binding, active storage in vacuoles, complexation by organic acids (Lee *et al.*, 1977, 1978; Mathys, 1977) and metal binding proteins and peptides, all play their part (Bradshaw *et al.*, 1990; Strange and Macnair, 1991; Shaw, 1999). There is also evidence that novel low-molecular weight compounds, the so-called phytochelatins (Salt *et al.*, 1989; Clemens *et al.*, 1999; Cobbett,

2000) and metallothioneins (Robinson, 1990; Robinson *et al.*, 1993; Zhou and Goldsbrough, 1994; Rauser, 1995; Zenk, 1996), may be involved in metal tolerance.

Although less than 1% of the Earth's land surface is covered by ultramafic rocks, the areas harbour a disproportionately higher percentage of endemic species. New Caledonia provides an exceptional case with 60% of the island's flora restricted to the ultramafics. Here, two monotypic families (Oncothecaceae and Strasburgeraceae), 38 genera (mostly monotypic!) and over 900 species are found on ultramafic soils (Kruckeberg and Rabinowitz, 1985; Jaffré *et al.*, 1987, 1997; Macnair and Gardner, 1998). Continental areas within South Africa, Zimbabwe, Brazil, Cuba (Brooks, 1987; Brooks *et al.*, 1990; Borhidi, 1995; Reeves *et al.*, 1999; Balkwill, 2001), and California also host a wide variety of species adapted to these extreme soils. In Cuba, 14.6% of the total flora (~920 taxa) is endemic, exclusively, to serpentine (Borhidi, 1995). This is especially remarkable given only 7% of the land surface of Cuba is of serpentinitic origin. In Zimbabwe, there are at least 20 species endemic to serpentine outcrops (Kruckeberg and Rabinowitz, 1985), while in California, 10% (215 species) of the flora endemic to that state is either wholly or largely restricted to serpentine soils (Kruckeberg, 1984).

The serpentine habitats of Sri Lanka: the background

The serpentine habitats in Sri Lanka have not been extensively studied; the authors are aware of only one published report by Brooks (1987). Brooks lists only three plant species (*Evolvulus alsinoides*, *Cymbopogon flexuosus*, *Morinda tinctoria*) in his brief account of the serpentines of the island indicating that the serpentine flora of Sri Lanka (as well as of India) is somewhat impoverished with regard to numbers of species as well as the percentage endemism. Brooks mentions that the floras do not hold the same potential for future botanical research as other Asian serpentine floras, particularly those of the offshore islands of Southeast Asia (the Malay Archipelago), where large numbers of endemic species have been recorded (Brooks, 1987; Proctor, 2003). Currently, there are no published records of any aspects of the fauna or specialized plant-insect or –microbial relationships (Hopkins, 1987; Boyd *et al.*, 1994; Boyd, 1998; Ghaderian *et al.*, 2000; Wall and Boyd, 2002; Héry *et al.*, 2003) of these sites, making the Sri Lankan sites some of the better documented geologically (Dissanayake and Van Riel, 1978; Munasinghe and Dissanayake, 1979, 1980; Dissanayake, 1982) yet natural historically least known serpentinized areas in Asia.

The observations made by Brooks (1987; personal communication, 1999) on Sri Lanka's serpentine flora during his brief visit to the island in 1981 are intriguing given the fact that Sri Lanka has greater biodiversity per unit area than any other country in Asia (Baldwin, 1991). Approximately 30% of the angiosperms (total 3100), 18% of the ferns (total 314), 35% of lichens (total 110), and 22% of the vertebrates (total 568) are endemic to the island. This high level of biological diversity and endemism, combined with the vulnerability of the habitats in which these species are often found, has attracted the attention of local and international conservation authorities. Sri Lanka, together with the Western Ghats of India, has now been declared one of 25 biodiversity hotspots in need of immediate protection (Myers *et al.*, 2000). Most of these conservation efforts and scientific studies are duly directed at the highly diverse lowland rainforests of the southwest where, for example, approximately 50% of the endemic angiosperms are locally restricted. The Sinharaja lowland rainforest, now a World Heritage Site with an area of 8,800 hectares or 89 km², is the home for 75, 95, 58, and 51% of Sri Lanka's endemic tree, bird, mammal, and butterfly species, respectively (Myers, 1990). While this level of diversity and endemism provides the need for special attention (De Zoysa and Raheem, 1990; Gunatilleke and Gunatilleke, 1985; Gunatilleke and Gunatilleke, 1991), it is also important that other habitat types, such as serpentine habitats, are also identified, characterized, and conserved. The lack of studies on the serpentine habitats of Sri Lanka indicates that there is generally a low awareness of the presence of this unusual habitat, with immense potential for harbouring unique assemblages of plants and animals.

In 1999, the first preliminary survey of four of the five known serpentine outcrops in Sri Lanka was conducted (Rajakaruna and Bohm,

2002). The survey was directed at describing the nature of the serpentine soils and exploring the floristic diversity found in these sites. To our knowledge, there have been no systematic studies previously done on the soils or flora of these sites. Thus, the paper added to the very limited knowledge on the serpentine soils and vegetation of the Indian subcontinent (Banerjee, 1972; Roy, 1974; Brooks, 1987) as well as that of Asia (Proctor, 2003). In the rest of this paper, we summarize the key findings reported in Rajakaruna and Bohm (2002) and a resulting pharmacological study using Sri Lanka's serpentine plants (Rajakaruna *et al.*, 2002b) in order to make our preliminary observations known to the larger scientific community in Sri Lanka. We hope that this paper will make regional scientists and conservationists aware of the unique nature of serpentine habitats and stimulate research as well as much-needed efforts to conserve these sites from any future destruction (Whiting *et al.*, 2002, 2004).

RESULTS AND DISCUSSION

Preliminary survey of Sri Lanka's serpentine habitats: a summary

There are at least five major serpentine outcrops in Sri Lanka (Figure 1). The following is a description of four of the sites explored during a brief visit to the island in 1999 (Rajakaruna and Bohm, 2002).

Site 1: Ussangoda

The Ussangoda (Welipatanwila) outcrop is located near Tangalla en route to Hambanthota on a coastal bluff overlooking the Indian Ocean. The area of the serpentine outcrop is approximately 3 km². The soil is characterized as hematite-rich clayey sand and is approximately 40 cm deep. The top 10 cm is overlain with very finely grained (loess-like) particles giving this heavily laterized site a reddish hue. This coloration has led to the common belief among local people that this is the site where betel, *Piper betel* L. (Piperaceae), was first brought to Sri Lanka - the red color of the soil resulting from the deposition of masticated waste of betel. Previous geochemical studies (Ranasinghe, 1987) have documented a total Ni content of 1000 ppm ($\mu\text{g/g}$

soil) and a Cr content of 7700 ppm ($\mu\text{g/g}$ soil) for these soils. The site is almost devoid of vegetation except for scattering of small herbaceous plants. The boundary of the serpentine is clearly marked by low thorn shrubs interspersed with a yet to be identified species of *Opuntia* (Cactaceae).

Site 2, 3: Indikolapelessa and Ginigalpelessa

The Indikolapelessa and Ginigalpelessa (Udawalawe) outcrops are located in the south-central part of the island. Each of these outcrops measures *c.* 1 km² in area. The soil is dark brown and poorly-developed. The total Ni content in the Ginigalpelessa site averages around 2100 $\mu\text{g/g}$, while the Cr content averages around 10,000 $\mu\text{g/g}$ (Ranasinghe, 1987). There is no geochemical information presently available for the site at Indikolapelessa. Dissanayake and Van Riel (1978, 1979) recorded a Ni content of 0.8-1.9% for the Udawalawe serpentinite.

Site 4: Yodhagannawa

The Yodhagannawa (Yudhaganawa) deposit is located in the north-central part of the island. There is presently no geochemical information available for the site. Interestingly, the reddish hue of the soil here has led to another folk belief that the red colour resulted from blood spilled during a battle between locals and Europeans during the colonial periods.

Both Ussangoda and Yodagannawa sites are protected. Ussangoda has been declared as a 'biologically sensitive area' while Yodagannawa is part of a nature reserve (Wasgamuwa National Park). While the latter may receive a higher level of protection, the site at Ussangoda is regularly used as a playground by local people and visitors. Flat portions of the outcrop are used as a cricket playing ground inflicting much damage to the soil as well as the limited number of herbaceous species that are present. Extensive use of this site will undoubtedly alter the soil properties and have a negative impact on the abundance of species present. The sites at Indikolapelessa and Ginigalpelessa are also somewhat protected owing to the fact that they are found within the boundaries of a state-run sugar cane plantation (Sevanagala Sugar Plantation). The staff indicated

that they have attempted to cultivate sugar cane in these sites but that the soil is not favourable for growth. While their lack of knowledge concerning the nature of serpentine substrate was apparent from these comments, it was also disturbing to realize that these sites may be used for future development, once it is realized that the sites are not suitable for cultivation. Both Ussangoda and Ginigalpelessa sites have been explored for geochemical prospects of various heavy metals, including gold (R. R. Brooks, R.

Ranasinghe, personal communications, 1999). Although it is known that high concentrations of both Ni and Cr have been found in these sites, the nature of the explorations or the decisions made regarding any future prospecting is unknown to the authors.




Figure 1. A map of Sri Lanka showing the approximate locations of the five serpentinite outcrops. Sites 1: Ussangoda; 2: Indikolapelessa; 3: Ginigalpelessa; 4: Yodhagannawa; 5: Katupotha (unexplored).

Serpentine soils of Sri Lanka

Since details of soil collection, sample preparation, and analyses are provided in Rajakaruna and Bohm (2002), we summarize the main findings for the purposes of this paper. The chemical and physical features of the soil samples collected from the serpentine sites in Sri Lanka are typical of such substrates found in other parts of the world (Brooks, 1987), with concentrations of DTPA-extractable (Lindsay and Norvell, 1978) heavy metals such as Ni and Cr uniformly high and Ca/Mg quotients consistently low. High heavy metal and low Ca concentrations make the soils unfavourable for normal plant growth. The

soils of Ussangoda differed the most from a typical serpentine soil. Here, pH and cation exchange capacity (CEC) were low and the Ca/Mg quotient was high. Further, Ussangoda soils are distinctive in having much higher concentrations of extractable Al, Co, and Zn and much lower concentrations of Fe, Mg, and Si. This may have resulted from heavy laterization of soils of this site, which results in loss of the divalent cations such as Fe, Mn, Co, and Ni through solubilization and leaching (Brooks, 1987). The unusually low CEC and low pH may also have contributed to loss of these ions as well as the greater availability of Al. Table 1 lists the

means for the various soil characteristics determined for these Sri Lankan sites.

Serpentine plants of Sri Lanka

The floristic survey documented a significantly larger number of species for the flora of Sri Lanka's serpentine habitats than previously recorded by Brooks (1987). For practical details on the methodology for plant collection, identification, and tissue analysis refer to Rajakaruna and Bohm (2002). The three dominant families represented are Poaceae, Asteraceae and Fabaceae (Table 2). The floristic composition was typical of serpentine habitats in other parts of the tropics with Asteraceae, Euphorbiaceae, Fabaceae, Poaceae and Rubiaceae well represented. All 21 families identified in the survey were known to occur in serpentine habitats elsewhere in the world. Further, at least 17 of the 40 genera identified are known from serpentine sites elsewhere (Table 3; Brooks, 1987, 1998; Baker *et al.*, 1992), suggesting that the survey has added perhaps 23 new genera to the list of plants that are known to occur on serpentine soils. Many of the genera (e.g., *Hybanthus*, *Evolvulus*), as well as the same species in some cases, occur in an arc-like distribution that runs between eastern Africa and Western Australia presenting an intriguing phytogeographic pattern. The total number of taxa collected per site were, Ginigalpelassa (25), Indikolapelassa (12), Ussangoda (9), and Yodhagannawa (3). Since the collections were made within a limited time span not all species growing in each site could be obtained. The most commonly found species, occurring in three of the four sites, were *Leucas zeylanica*, *Cymbopogon flexuosus*, *Eupatorium odoratum*, and *Phoenix farinifera* all found in two sites each. Surprisingly, none of the species identified was endemic to the island. However, the precise taxonomy of several species is questionable. These include: *Hybanthus enneaspermus*, *Evolvulus alsinoides*, *Crotalaria biflora*, *Desmodium triflorum*, *Epaltes divaricata*, *Geniosporum tenuiflorum*, *Fimbristylis falcata*, *Cymbopogon flexuosus* as well as an unidentified species each of *Canthium* and *Phyllanthus*. One of the more interesting observations made was in the colour polymorphism of *Evolvulus alsinoides*. Whilst it is known that the species has two colour morphs (Austin in Dassanayake and Fosberg, 1980), it is unknown whether there is a correlation between colour and any edaphic

differences in their microhabitats. Such edaphically-correlated pigment differences are known in species that inhabit distinct soils (Rajakaruna and Bohm, 1999; Schemske and Bierzychudek, 2001). In comparison to serpentine floras of the islands of southeast Asia, the Sri Lankan serpentine flora appears to be impoverished with regard to both number of species as well as degree of endemism. It is unclear why an island as rich in endemics as Sri Lanka should harbour such an apparently impoverished serpentine flora. A factor that may have contributed to this apparent anomaly may be that sampling in this survey was limited to those species that could be identified using floral or other reproductive structures. Rare or endemic taxa could have been easily overlooked if they were not in flowering condition. Further, sampling was carried out over only a day or even hours depending on the site, seriously limiting opportunities for a thorough survey of the flora. Limited sampling also prevented observation of any minor morphological features that might indicate taxonomically recognizable differences especially in those taxa mentioned earlier where there were obvious differences in plant size and growth habit relative to populations that are found off the serpentine substrate (personal observations). For example, at least 17 varieties of *Evolvulus alsinoides* have been described globally (Austin in Dassanayake and Fosberg, 1980) and the small-leaved, prostrate growth habit plants found at Ussangoda may represent an edaphic race or even subspecies restricted to the serpentine substrate. Brooks (1987) states that *Cymbopogon flexuosus* found in the Sri Lankan serpentine sites is a form endemic to serpentine. This observation made by Brooks (1987) is worthy of further investigation.

Tissue analyses indicate that elemental concentrations found in the Sri Lankan serpentine plants (Rajakaruna and Bohm, 2002) are typical of those found in serpentine plants elsewhere (Brooks, 1987, 1998; Roberts and Proctor, 1992; Baker *et al.*, 1992). Most of the plants collected accumulated higher than 'normal' concentrations of several elements: Al, As, Cd, Co, Cu, Cr, Na, Ni, Pb, and Zn. Concentrations of all these elements were greater than those that are normally considered to be toxic to most plants (Kabata-Pendias and Pendias, 1984; Mehta and Farago, 1994). Of particular interest was the discovery of

three Ni and five Cu hyperaccumulators (Table 4). Hyperaccumulators demonstrate extreme metal tolerance and are able to accumulate over 100 times the concentration found in non-hyperaccumulator plants growing on the same substrate (Brooks, 1998). For most metals, hyperaccumulation is defined as more than 1000 µg metal per g dry leaf tissue (0.1%) however for Zn and Mn, 1% is regarded as exceptional (Baker and Brooks, 1989). All three Ni hyperaccumulators were discovered at Ussangoda. Two of the three Ni hyperaccumulators discovered, *Evolvulus alsinoides* and *Hybanthus enneaspermus*, also occur in serpentine sites in Queensland, Australia (R. D. Reeves, Massey University, New Zealand, personal communication), but neither of these accumulates sufficient Ni to qualify as a hyperaccumulator. Since *E. alsinoides* is known to comprise numerous varieties, it is tempting to speculate that the population found at Ussangoda may represent a physiologically distinct form, perhaps an edaphic race. At least seven taxa from the genus *Hybanthus* are known to hyperaccumulate Ni (Brooks, 1998; Reeves, 1992, 2003; Reeves and Baker, 2000). Given the predisposition of taxa from this genus to hyperaccumulate Ni, it is possible that the population found at Ussangoda may possess this unique capacity. Both species, however, warrant further detailed investigation to confirm these preliminary findings. The third Ni hyperaccumulator revealed by the survey, *Crotalaria biflora*, has not been previously recorded to hyperaccumulate this metal in other parts of its range. However, *C. cobalticola* Duvign. & Plancke from the Democratic Republic of Congo (formerly Zaïre), is a known Co hyperaccumulator (Brooks *et al.*, 1995). It would be of interest to find out if the tendency to hyperaccumulate metals is more widespread in this large genus.

Rinorea bengalensis (Wall.) O.K. (Violaceae to which *Hybanthus* also belongs), a species that was not encountered in the current survey, also hyperaccumulates Ni (Brooks and Wither 1977a, b; Jopony and Baker, 2000).

Rinorea bengalensis is found in a broad arc-like distribution encompassing Sri Lanka, the Malay Archipelago, New Guinea, the Solomon Islands, and Queensland, Australia. Brooks and Wither (1977a, b) conducted a survey of herbarium specimens from the entire range of this species and found that Ni hyperaccumulation is widespread in this plant. The level of Ni found in the herbarium specimen of this species from Sri Lanka was reported to be in excess of 10,000 ppm (10-fold the minimum required to be a hyperaccumulator!). This high level indicates that the specimen tested was obviously collected from a serpentine site, the locality indicated on a map presented by Brooks and Wither (1977a) suggesting a collection in the central part of the island. This species was not encountered in our field exploration and is worthy of intense exploration, especially in the Udawalawe sites or at the unexplored site near Katupotha.

A surprising finding of the tissue analyses was the discovery of five Cu hyperaccumulators: *Geniosporum tenuiflorum*, *Clerodendrum infortunatum*, *Croton bonplandianus*, *Waltheria indica*, and *Tephrosia villosa* all hyperaccumulated this metal. To date, almost all Cu hyperaccumulators discovered have been from the copper mineralized soils of Democratic Republic of Congo (Brooks *et al.*, 1992; Reeves and Baker, 2000). Hyperaccumulation of Cu by serpentine plants appears to be a very rare phenomenon. However, tolerance to potentially toxic levels of Cu has been observed in *Mimulus guttatus* Fischer ex DC (Scrophulariaceae), a species that occurs in serpentine soils in California (Tilstone and Macnair, 1997a, b). Although none of the Cu hyperaccumulators discovered in Sri Lanka has been previously known to have this capacity, both Fabaceae and Lamiaceae (three of the five taxa belong to these two families) have genera with the capacity to hyperaccumulate Cu (Reeves and Baker, 2000). The discovery of these new Cu hyperaccumulators from the ultramafics of Sri Lanka is even more surprising since Brooks in a biogeochemical and geobotanical survey of the Seruwila copper-magnetite prospect near

Table 1.

The species collected from the four serpentine sites in Sri Lanka. Site abbreviations are GGP=Ginigalpelassa, IP=Indikolapelassa, U=Ussangoda, Y=Yodhagannawa.

Family	Species	Site
Asteraceae	<i>Epaltes divaricata</i> (L.) Cass.	U
Convolvulaceae	<i>Evolvulus alsinoides</i> (L.) L.	U
Cyperaceae	<i>Fimbristylis falcata</i> (Vahl.) Kunth	U
Fabaceae	<i>Cassia auriculata</i> L.	U
Fabaceae	<i>Crotalaria biflora</i> (L.) L.	U
Fabaceae	<i>Desmodium triflorum</i> (L.) DC.	U
Rubiaceae	<i>Tarenna asiatica</i> (L.) Alston	U
Rutaceae	<i>Toddalia asiatica</i> Lamk.	U
Violaceae	<i>Hybanthus enneaspermus</i> (L.) F. Muell.	U
Fabaceae	<i>Tephrosia purpurea</i> Pers.	Y
Lamiaceae	<i>Geniosporum tenuiflorum</i> (L.) Merr.	Y
Lamiaceae	<i>Leucas zeylanica</i> (L.) R. Br.	Y
Arecaceae	<i>Phoenix farinifera</i> Roxb.	IP
Asteraceae	<i>Eclipta prostrata</i> (L.) L.	IP
Asteraceae	<i>Eupatorium odoratum</i> L.	IP
Asteraceae	<i>Spilanthes calva</i> DC.	IP
Asteraceae	<i>Vernonia cinerea</i> (L.) Less.	IP
Euphorbiaceae	<i>Phyllanthus myrtifolius</i> Moon	IP
Fabaceae	<i>Tephrosia villosa</i> Pers.	IP
Lamiaceae	<i>Leucas zeylanica</i> (L.) R. Br.	IP
Malvaceae	<i>Abutilon indicum</i> Sweet	IP
Malvaceae	<i>Sida acuta</i> Burm.	IP
Poaceae	<i>Cymbopogon flexuosus</i> (Steudel) W. Watson.	IP
Solanaceae	<i>Physalis minima</i> L.	IP
Amaranthaceae	<i>Aerva lanata</i> (L.) Juss. Ex Schult.	GGP
Apocynaceae	<i>Carissa spinarum</i> L.	GGP
Arecaceae	<i>Phoenix farinifera</i> Roxb.	GGP
Asclepiadaceae	<i>Calotropis gigantea</i> (L.) R. Br.	GGP
Asteraceae	<i>Eupatorium odoratum</i> L.	GGP
Euphorbiaceae	<i>Croton bonplandianus</i> Baill.	GGP
Euphorbiaceae	<i>Croton hirtus</i> L'Her.	GGP
Euphorbiaceae	<i>Euphorbia rubicunda</i> Blume	GGP
Euphorbiaceae	<i>Phyllanthus</i> L. sp.	GGP
Fabaceae	<i>Cassia fistula</i> L.	GGP
Lamiaceae	<i>Leucas zeylanica</i> (L.) R. Br.	GGP
Lamiaceae	<i>Ocimum sanctum</i> L.	GGP
Liliaceae	<i>Asparagus zeylanicus</i> Hook. f.	GGP
Myrtaceae	<i>Syzygium cumini</i> Skeels	GGP
Poaceae	<i>Apluda mutica</i> L.	GGP
Poaceae	<i>Aristida setacea</i> Retz.	GGP
Poaceae	<i>Eragrostis ciliaris</i> (L.) R. Br.	GGP
Poaceae	<i>Cymbopogon flexuosus</i> (Steudel) W. Watson	GGP
Rhamnaceae	<i>Zizyphus oenoplia</i> Mill.	GGP
Rubiaceae	<i>Canthium puberulum</i> Thw. ex. Hook. f.	GGP
Rubiaceae	<i>Canthium</i> Lam. sp.	GGP
Rubiaceae	<i>Morinda tinctoria</i> Roxb.	GGP
Sterculiaceae	<i>Waltheria indica</i> L.	GGP
Verbenaceae	<i>Clerodendrum infortunatum</i> L.	GGP
Verbenaceae	<i>Lantana camara</i> L.	GGP

Trincomalee did not find any plant with a copper concentration $> 21 \mu\text{g/g}$ in leaf dry matter.

The occurrence of hyperaccumulators in specific families and within certain genera in various parts of the world raises the question as to whether some families as well as genera possess a constitutive or biochemical factor that predisposes them to serpentine tolerance accompanied by the accumulation of large amounts of metal (Bradshaw *et al.*, 1990). So far >320 (and growing!) hyperaccumulators of Ni have been discovered (Reeves and Baker, 2000).

Brassicaceae and Euphorbiaceae include 82 species each while Asteraceae, Flacourtiaceae, Buxaceae, and Rubiaceae include 26, 19, 17, and 12 species, respectively (Reeves and Baker, 2000; Borhidi, 2001). Of the 24 or so known Cu hyperaccumulators, 14 belong to Cyperaceae, Lamiaceae, Poaceae, Scrophulariaceae, Fabaceae and Asteraceae (Baker and Brooks, 1989; Reeves and Baker, 2000). The hyperaccumulators of Ni and Cu discovered in Sri Lanka add to these lists further confirming the observation that certain families are apparently predisposed to having species with the capacity to accumulate metals.

Table 2.

Mean (\pm SE) for various soil features in soil samples collected from the four ultramafic sites in Sri Lanka. Element concentrations are in $\mu\text{g/g}$ dry soil and CEC is in $\text{Cmol}^{(+)}/\text{kg}$ dry soil. The mean concentrations for the four sites (L-R) are a result of soil analysis of 5, 6, 6, and 3 samples, respectively.

Soil Feature	Ginigalpelessa	Indikolapelessa	Ussangoda	Yodhagannawa
Aluminum	0.2 (\pm 0.07)	0.5 (\pm 0.2)	6.6 (\pm 1)	0.13 (\pm .02)
Barium	1(\pm 0.52)	0.97 (\pm 0.21)	0.3 (\pm 0.08)	0.2 (\pm 0.01)
Boron	0.01 (\pm 0.006)	0.01 (\pm 0.006)	0.04 (\pm 0.002)	0.01 (\pm 0.01)
Cadmium	0.0002 (\pm 0.0002)	0.02 (\pm 0.01)	0.002 (\pm 0.002)	0
Calcium	921 (\pm 294)	1030 (\pm 239)	166 (\pm 13.9)	128 (\pm 4.6)
Ca/Mg	0.35 (\pm 0.1)	1.04 (\pm 0.5)	1.83 (\pm 0.2)	0.14 (0.01)
CEC	42.2 (\pm 3.6)	34.3 (\pm 3.6)	8.5 (\pm 0.8)	15.4 (\pm 0.7)
Chromium	0.07 (\pm 0.03)	0.05 (\pm 0.01)	0.05 (\pm 0.01)	0.05 (\pm 0.01)
Cobalt	1.94 (\pm 0.56)	2.56 (\pm 0.94)	5.3 (\pm 0.98)	1.4 (\pm 0.14)
Copper	0.75 (\pm 0.15)	1.4 (\pm 0.6)	0.62 (\pm 0.05)	0.22 (\pm 0.02)
Iron	58.2 (\pm 10.6)	39.9 (\pm 4.5)	16.5 (\pm 0.6)	46.8 (\pm 6.6)
Lead	0.15 (\pm 0.03)	0.13 (\pm 0.04)	0.25 (\pm 0.06)	0.05 (\pm 0.01)
Magnesium	2810 (\pm 164.1)	1752.1(\pm 357.8)	94.6 (\pm 10.4)	937.5 (\pm 50.5)
Manganese	32.8 (\pm 10.3)	76.1(\pm 18.3)	63.8 (\pm 8.2)	25.7 (\pm 2.4)
Molybdenum	0.001(\pm 0.0006)	0.002 (\pm 0.0017)	0	0
Nickel	104.2 (\pm 37.1)	59.2 (\pm 21.2)	45.6 (\pm 5.9)	66.9 (\pm 9.9)
pH (water)	6.3 (\pm 0.3)	5.6 (\pm 0.2)	4.6 (\pm 0.1)	5.4 (\pm 0.2)
pH (CaCl ₂)	5.7 (\pm 0.4)	5.3 (\pm 0.2)	4.3 (\pm 0.1)	5 (\pm 0.2)
Potassium	144 (\pm 29.5)	419.6 (\pm 232.4)	224.2 (\pm 27.9)	67.5 (\pm 7.5)
K/Na	12.8 (\pm 5)	39.1(\pm 15)	12.5 (\pm 2.9)	8.3 (\pm 0.8)
Silicon	13.1(\pm 1.6)	19.6 (\pm 5.4)	0.4 (\pm 0.05)	18 (\pm 0.9)
Sodium	26.7 (11.4)	8.9 (\pm 1.1)	23.2 (\pm 6.7)	8.1 (\pm 0.5)
Zinc	0.63 (\pm 0.2)	0.92 (\pm 0.2)	2.1(\pm 1.4)	0.7 (\pm 0.08)

Table 3.

The phytogeography and previously documented metal hyperaccumulating behaviors of 17 genera found on the serpentine substrate in Sri Lanka. Information from various sources listed in Brooks, 1987, 1998; Baker *et al.*, 1992; Reeves and Baker, 2000 and Reeves, personal communication, 2000.

Genus (Family)	Recorded Distribution on Serpentine	Metal Hyperaccumulating Behavior (if documented)
<i>Aristida</i> (Poaceae)	Zimbabwe	
<i>Cassia</i> (Fabaceae)	Australia	
<i>Crotalaria</i> (Fabaceae)	Angola, Zimbabwe, Zaire	Cobalt (1 taxon)
<i>Cymbopogon</i> (Poaceae)	Oman, Zimbabwe	
<i>Desmodium</i> (Fabaceae)	Maryland, U.S.A.	
<i>Epaltes</i> (Asteraceae)	W. Australia	
<i>Eragrostis</i> (Poaceae)	Zimbabwe	
<i>Eupatorium</i> (Asteraceae)	Maryland, U.S.A.	
<i>Euphorbia</i> (Euphorbiaceae)	Anatolia, Brazil, Oman, Zimbabwe, Cuba	Nickel (2 taxa)
<i>Evolvulus</i> (Convolvulaceae)	W. Australia	
<i>Fimbristylis</i> (Cyperaceae)	South Africa, New Caledonia, W. Australia	
<i>Hybanthus</i> (Violaceae)	New Caledonia, W. Australia, Cuba	Nickel (7 taxa)
<i>Leucas</i> (Lamiaceae)	Zimbabwe	
<i>Morinda</i> (Rubiaceae)	Philippines	
<i>Phyllanthus</i> (Euphorbiaceae)	Philippines, Sabah, New Caledonia, Cuba	Nickel (10 taxa)
<i>Syzygium</i> (Myrtaceae)	Philippines	
<i>Tarenna</i> (Rubiaceae)	New Caledonia	

Table 4.

A sample of species showing high metal accumulating behaviour collected from the Sri Lankan serpentine sites (from Rajakaruna and Bohm, 2002). Soil concentrations, when listed, are *extractable* or *exchangeable* values determined for soils collected from 0-10cm depth. Plant tissue concentrations are *total* values for whole plants in herbaceous species and shoots in shrubs. Hyperaccumulators of Ni and Cu are in bold. Site abbreviations are GGP=Ginigalpelassa, IP=Indikolapelassa, U=Ussangoda, Y=Yodhagannawa.

Species	Site	Element	Soil ($\mu\text{g/g}$ dry)	Plant ($\mu\text{g/g}$ dry)
<i>Aristida setacea</i>	GGP	Cr	0.1	19
		Ni	15	439
<i>Calotropis gigantea</i>	GGP	Na	66	12017
<i>Canthium sp.</i>	GGP	Cr	-	16
<i>Cassia auriculata</i>	U	Cu	0.5	885
<i>Clerodendrum infortunatum</i>	GGP	Cu	-	2208
<i>Crotalaria biflora</i>	U	Ni	42	1048
<i>Croton bonplandianus</i>	GGP	Cu	-	2206
<i>Croton hirtus</i>	GGP	Cr	-	40
<i>Cymbopogon flexuosus</i>	GGP	Cr	-	26
	IP	Cr	-	18
<i>Epaltes divaricata</i>	U	Ni	52	733
		Cr	0.1	17
<i>Evolvulus alsinoides</i>	U	Ni	55	1297
		Cr	0.1	23
<i>Geniosporum tenuiflorum</i>	Y	Cr	0.1	21
		Cu	0.3	2299.3
<i>Hybanthus enneaspermus</i>	U	Ni	29	1786
		Cr	0.1	21
<i>Leucas zeylanica</i>	GGP	Cr	0.1	28
	IP	Cr	0.1	19
<i>Phyllanthus sp.</i>	GGP	Cu	-	821
<i>Phyllanthus myrtifolius</i>	IP	Cr	-	22
<i>Tephrosia villosa</i>	IP	Cu	0.2	1723
<i>Waltheria indica</i>	GGP	Cu	-	1534

CONCLUSIONS

The study reported in Rajakaruna and Bohm (2002) and summarized here represents the

first broad-scale floristic survey of the serpentine habitats of Sri Lanka. Owing to time constraints, sampling was limited and was restricted to those

species identifiable using reproductive structures. This type of haphazard sampling may have had an effect on the unexpected paucity of endemic or rare species. Reiterating the preliminary nature of this survey, it should also be pointed out that no attempts were made to determine cellular localizations of any of the accumulated metals or investigate any other physiological aspects of the plant species found at these sites. The findings, however, warrant detailed collections, expanded taxonomic study, and potentially significant ecophysiological investigations of Ni and Cu hyperaccumulation.

Serpentine habitats provide a multitude of opportunities for botanical research. In addition to taxonomic and physiological studies of the nature described in this paper, serpentine habitats also provide unique opportunities to explore specialized plant-microbe interactions (Hopkins, 1987; Ghaderian *et al.*, 2000; Hungate *et al.*, 2000; Jaffré *et al.*, 2001) and plant-herbivore interactions (Boyd, 1998; Mesjasz-Przybylowicz and Przybylowicz, 2001), the process of speciation (Kruckeberg, 1996; Macnair and Baker, 1994; Macnair and Gardner, 1998; Rajakaruna and Whitton, 2004), as well discovery of novel compounds that may be produced in response to edaphic stresses. It is possible such compounds could have secondary and often unforeseen effects on other organisms (Hanawa *et al.*, 2000; Pernas *et al.*, 2000). In fact, a study conducted on the pharmacological properties of the Sri Lankan serpentine plants (Rajakaruna *et al.*, 2002b) document that 29 of the 45 species are active against at least one microorganism tested, suggesting that these plants may produce chemical compounds with important antimicrobial properties.

Serpentine habitats of Sri Lanka are easily accessible and can provide a living laboratory for numerous scientific studies both exploratory and experimental in nature. In many ways, they can provide a counterpart to the highly regarded and carefully studied rainforests of the southwest. The lack of information and awareness of the serpentines of Sri Lanka will only lead to the destruction of these unique habitats that have so far gone unnoticed as potential sites for harbouring a wealth of genetic diversity (Whiting *et al.*, 2002, 2004).

ACKNOWLEDGEMENTS

We thank Dr. Ananda Gunatilleke for directing the attention of the primary author to the serpentine habitats of Sri Lanka; Mr. Nimal Ranasinghe, Geological Survey Department for providing site information and sharing unpublished soil data; Professor Emeritus M. D. Dassanayake and Ms. R. W. Edirisinghe, National Herbarium, Peradeniya for assistance in plant identification; Staff, Sevanagala Sugar Plantation for allowing access to sites and helping in the sampling; Dr. Wasantha Seneviratne, Dept. of Botany, Univ. of Colombo for sharing her interest on the serpentines; Drs. Sarath and Jayanthi Edirisinghe for assisting in field collections; Department of Wildlife Conservation for granting permission to export soil and tissue samples for the purposes of study; Dr. Nimal Gunatilleke, Dept. of Botany, Univ. of Peradeniya for encouraging the writing of this paper.

REFERENCES

- Antonovics, J., Bradshaw, A. D. and Turner, R. G. (1971). Heavy metal tolerance in plants. *Advances in Ecological Research* **7**: 1-85.
- Arianoutsou, M., Rundel, P. W. and Berry, W. L. (1993). Serpentine endemics as biological indicators of soil elemental concentrations. In: *Plants as Biomonitors. Indicators for Heavy Metals in the Terrestrial Environment* (Ed. B. Markert), VCH, Weinheim, Germany.
- Austin, D. F. (1980). Convolvulaceae. In: *A Revised Handbook to the Flora of Ceylon*. Vol.I. (Eds. M. D. Dassanayake and F. R. Fosberg), pp. 288-363. A. A. Balkema, Rotterdam, The Netherlands.
- Baker, A. J. M. (1987). Metal tolerance. In: *Frontiers of Comparative Plant Ecology*, (Ed. by I. H. Rorison, J. P. Grime, R. Hunt, G. A. F. Hendry and D. H. Lewis), Academic Press, New York & London, and *New Phytologist* **106** (Suppl.): 93-111.
- Baker, A. J. M. and Brooks, R. R. (1988). Botanical exploration for minerals in the humid tropics. *Journal of Biogeography* **15**: 221-229.
- Baker, A. J. M. and Brooks, R. R. (1989).

- Terrestrial higher plants which hyperaccumulate metallic elements - a review of their distribution, ecology and phytochemistry. *Biorecovery* **1**: 81-126.
- Baker, A. J. M., McGrath, S. P., Reeves, R. D. and Smith, J. A. C. (2000). Metal hyperaccumulator plants: a review of the ecology and physiology of a biological resource for phytoremediation of metal-polluted soils. In: *Phytoremediation of Contaminated Soil and Water*, (Ed. by N. Terry and G. S. Bañuelos), pp. 85-107. CRC Press Inc, Boca Raton, FL, USA.
- Baker, A. J. M., Proctor J. and Reeves R. D. (Eds). (1992). *The Vegetation of Ultramafic (Serpentine) Soils*. Proceedings of the First International Conference on Serpentine Ecology, University of California, Davis, 19-22 June 1991. Intercept, Hampshire, UK. 509 p.
- Baker, A. J. M. and Walker, P. L. (1990). Ecophysiology of metal uptake by tolerant plants. In: *Heavy Metal Tolerance in Plants: Evolutionary Aspects* (Ed. A. J. Shaw), pp. 155-177. CRC Press, Boca Raton, FL, USA.
- Baldwin, M. F. (Ed.) (1991). *Natural Resources of Sri Lanka. Conditions and Trends*. A Report Prepared for the Natural Resources, Energy and Science Authority of Sri Lanka. Keels Business Systems Ltd., Colombo, Sri Lanka.
- Balkwill, K. (Ed.) (2001). Proceedings of the Third International Conference on Serpentine Ecology, Berg-en-Dal, Kruger National Park, South Africa, 22-28 March 1999 *South African Journal of Science* **97**, 485-628.
- Banerjee, P. K. (1972). Geology and geochemistry of the Sukinda ultramafic field, Cuttack District, Orissa. *Memoirs of the Geological Survey of India* **103**: 171.
- Black, C. A., Evans D. D., White J. L., Ensminger L. E. and Clark F. E. (1965). *Methods of Soil Analysis. Chemical and Microbial Properties*. Part 2. American Society of Agronomy Inc., Madison, WI, USA.
- Borhidi, A. (1995). *Phytogeography and Vegetation Ecology of Cuba*. Akademiai-Kiado, Budapest, Hungary. 752 p.
- Borhidi, A. (2001). Phylogenetic trends in Ni-accumulating plants. *South African Journal of Science* **97**: 544-547.
- Boyd, R. S. and Martens, S. N. (1992). The raison d'être for metal hyperaccumulation by plants. In: *The Vegetation of Ultramafic (Serpentine) Soils*. Proceedings of the First International Conference on Serpentine Ecology (Eds. A. J. M. Baker, J. Proctor and R. D. Reeves), pp. 279-289. Intercept, Hampshire, UK.
- Boyd, R. S., Shaw, J., Martens, S. N. (1994). Nickel hyperaccumulation defends *Streptanthus polygaloides* (Brassicaceae) against pathogens. *American Journal of Botany* **81**: 294-300.
- Boyd, R. S. and Martens, S. N. (1998). The significance of metal hyperaccumulation for biotic interactions. *Chemoecology* **8**: 1-7.
- Bradshaw, A. D., McNeilly, T. and Putwain, P. D. (1990). The essential qualities. In: *Heavy Metal Tolerance in Plants: Evolutionary Aspects* (Ed. A. J. Shaw), pp. 323-334. CRC Press, Boca Raton, FL, USA.
- Brooks, R. R. and Wither, E. D. (1977a). Nickel hyperaccumulation by *Rinorea bengalensis* (Wall.) O.K.. *Journal of Geochemical Exploration* **7**: 295-300.
- Brooks, R. R. and Wither, E. D. (1977b). Cobalt and nickel in *Rinorea* species. *Plant and Soil* **47**: 707-712.
- Brooks, R. R. (1972). *Geobotany and Biogeochemistry in Mineral Exploration*. Harper and Row, New York, USA, 290 p.
- Brooks, R. R. (1987). *Serpentine and its Vegetation. A Multidisciplinary Approach*. Dioscorides Press, OR, USA, 322 p.
- Brooks, R. R. and Johannes, D. (1990). *Phytoarchaeology*. Dioscorides Press, Portland, OR, USA, 224 p.

- Brooks, R. R. (1994). Plants that hyperaccumulate heavy metals. In: *Plants and the Chemical Elements* (Ed. M. E. Farago), pp. 88-105. VCH, Weinheim, Germany.
- Brooks, R. R., Dunn, C. E. and Hall G. E. M. (1995). *Biological Systems in Mineral Exploration and Processing*. Ellis Horwood, Hemel Hempstead, UK. 539 p.
- Brooks, R. R. (1998). *Plants that Hyperaccumulate Heavy Metals*. CAB International, Wallingford, UK. 380 p.
- Brooks, R. R., Reeves, R. D., Baker, A. J. M., Rizzo, J. A. and Diaz Ferreira, H. (1990). The Brazilian Serpentine Plant Expedition (BRASPEX), 1988. *National Geographic Research* **6**: 205-219.
- Brooks, R. R., Baker, A. J. M. and Malaisse, F. (1992). Copper flowers: the unique flora of the copper hills of Zaïre. *National Geographic Research & Exploration* **8**: 338-351.
- Brooks, R. R., Baker, A. J. M., Ramakrishna, R. S. and Ryan, D. E. (1985). Botanical and geochemical exploration studies at the Seruwila copper-magnetite prospect in Sri Lanka. *Journal of Geochemical Exploration* **24**: 223-235.
- Brooks, R. R., Chambers, M. F., Nicks, L. J. and Robinson, B. H. (1998). Phytomining. *Trends in Plant Science* **3**: 359-362.
- Cannon, H. L. (1960). Botanical prospecting for ore deposits. *Science* **132**: 591-598.
- Cannon, H. L. (1971). The use of plant indicators in ground water surveys, geologic mapping, and mineral prospecting. *Taxon* **20**: 227-256.
- Clemens, S., Kim, E. J., Neumann, D. and Schroeder, J. I. (1999). Tolerance to toxic metals by a gene family of phytochelatin synthases from plants and yeast. *European Molecular Biology Organization Journal* **18**: 3325-3333.
- Cobbett, C. S. (2000). Phytochelatins and their roles in heavy metal detoxification. *Plant Physiology* **123**: 825-832.
- Coleman, R. G. and Jove, C. (1992). Geological origin of serpentinites. In: *The Vegetation of Ultramafic (Serpentine) Soils*. Proceedings of the First International Conference on Serpentine Ecology (Eds A. J. M. Baker, J. Proctor and R. D. Reeves), pp. 1-17. Intercept, Hampshire, UK.
- De Zoysa, N. and Raheem, R. (1990). *Sinharaja. A rainforest in Sri Lanka*. Aitken Spence and Co. Ltd., Colombo, Sri Lanka. 63 p.
- Dissanayake, C. B. and Van Riel, B. J. (1978). Petrology and geochemistry of a recently discovered nickeliferous serpentinite from Sri Lanka. *Journal of the Geological Society of India* **19**, 464-471.
- Dissanayake, C. B. and Van Riel, B. J. (1979). Petrology and geochemistry of a recently discovered nickeliferous serpentinite from Sri Lanka: authors' reply. *Journal of the Geological Society of India* **20**, 302.
- Dissanayake, C. B. (1982). Recognition of ultramafic rocks and nickeliferous laterites in humid tropical terrains: some guidelines. In: *Hidden Wealth: Mineral Exploration Techniques in Tropical Forest Areas*. Proceedings/Symposium on Mineral Exploration Techniques in Tropical Forest Areas (Eds D. J. C. Laming and A. K. Gibbs). Association of Geoscientists for International Development, Asian Institute of Technology.
- Farago, M. E. and Cole, M. M. (1988) Nickel and plants. In: *Metal ions in biological systems. Volume 23. Nickel and its Role in Biology* (Eds. H. Sigel and A. Sigel), Marcel Dekker Inc., New York, USA.
- Gabbrielli, R., Mattioni, C. and Vergnano, O. (1991). Accumulation mechanisms and heavy metal tolerance of a nickel hyperaccumulator. *Journal of Plant Nutrition* **14**: 1067-1080.
- Ghaderian, Y. S. M., Lyon, A. J. E., Baker, A. J. M. (2000). Seedling mortality of metal hyperaccumulator plants resulting from damping off by *Pythium* spp. *New Phytologist* **146**: 219-224.

- Gordon, A. and Lipman, C. B. (1926). Why are serpentine and other magnesian soils infertile? *Soil Science* **22**: 291-302.
- Gunatilleke, C. V. S. and Gunatilleke, I. A. U. N. (1985). Phytosociology of Sinharaja – A contribution to rainforest conservation in Sri Lanka. *Biological Conservation* **31**: 21-40.
- Gunatilleke, I. A. U. N. and Gunatilleke, C. V. S. (1991). Threatened woody endemics of the wet lowlands of Sri Lanka and their conservation. *Biological Conservation* **55**: 17-36.
- Hanawa, F., Tahara, S., Towers, G. H. N. (2000). Antifungal nitro compounds from skunk cabbage (*Lysichitum americanum*) leaves treated with cupric chloride. *Phytochemistry* **53**: 55-58.
- Héry, M., Nazaret, S., Jaffré, T., Normand, P., and E. Navarro. (2003). Adaptation to nickel spiking of bacterial communities in neocaledonian soils. *Environmental Microbiology* **5**: 3-12.
- Hopkins, N. A. (1987). Mycorrhizae in a California serpentine grassland community. *Canadian Journal of Botany* **65**: 484-487.
- Hungate, B. A., Jaeger, C. H., Gamara, G., Chapin, F. S., Field, C.B. (2000). Soil microbiota in two annual grasslands: responses to elevated atmospheric CO₂. *Oecologia* **124**: 589-598.
- Jenny, H. (1980). *The Soil Resource. Origin and Behaviour*. Springer-Verlag. New York., USA, 377 p.
- Jaffré, T., Morat, P., Veillon, J. M. and MacKee, H. S. (1987). Changements dans la végétation de la Nouvelle Calédonie ou cours du Tertiaire: la végétation et la flora des roches ultrabasiqes. *Adansonia* **4**: 365-391.
- Jaffré, T., Reeves, R. D., Becquer, T. (Eds.). (1997). *The Ecology of Ultramafic and Metalliferous Areas*. Proceedings of the Second International Conference on Serpentine Ecology, Nouméa, New Caledonia, 31 July – 5 August, 1995. Documents Scientifiques et Techniques, No. III2. Nouméa, New Caledonia: Centre ORSTOM de Nouméa, 306 p.
- Jaffré, T., McCoy, S. G., Rigault, F., Navarro, E. (2001). A comparative study of flora and symbiotic micro-flora diversity in two *Gymnostoma* formations of ultramafic rocks in New Caledonia. *South African Journal of Science* **97**: 599-603.
- Jopony, M. and Baker, A. J. M. (2000). Hyperaccumulation of nickel and zinc by plants of Southeast Asia. In: Proceedings of the SoilRem 2000 Conference, Hangzhou, PR China, 15-19 October 2000 (Ed. by Y. M. Luo, Z. H. Cao, Y. X. Chen, S. P McGrath, F. J. Zhao and J. M. Xu) pp. 188-190.
- Kabata-Pendias, A. and Pendias, H. (1984). *Trace Elements in Soils and Plants*. CRC Press, Boca Raton, FL, USA, 315 p.
- Krause, W. (1958). *Andere Bodenspezialisten. Handb. Pflanzenphysiol.* **4**: 755-806.
- Kruckeberg, A. R. (1984). *California Serpentine: Flora, Vegetation, Geology, Soils, and Management Problems*. University of California Press. Berkeley, California, USA 180 p.
- Kruckeberg, A. R. and Rabinowitz, D. (1985). Biological aspects of endemism in higher plants. *Annual Reviews of Ecology and Systematics* **16**: 447-479.
- Kruckeberg, A. R. (1986). An essay: The stimulus of unusual geologies for plant speciation. *Systematic Botany* **11**: 455-463.
- Kruckeberg, A. R. (1992). Plant life of Western North American ultramafics. In: *The Ecology of Areas with Serpentinized Rocks. A World View* (Eds. B. A. Roberts and J. Proctor), pp. 31-73. Kluwer Academic Publications, Dordrecht, The Netherlands.
- Kruckeberg, A. R., Peterson, P. J. and Semiullah, Y. (1993). Hyperaccumulation of nickel by *Arenaria rubella* (Caryophyllaceae) from Washington Sate. *Madroño* **40**: 25-30.
- Kruckeberg, A. R. (2002). *Geology and Plant Life. The Effects of Landforms and Rock Types on Plants*. University of Washington Press, Seattle, USA. 362 p.

- Lee, J., Reeves, R. D., Brooks, R. R. and Jaffré, T. (1977). Isolation and identification of a citrato complex of nickel from nickel-accumulating plants. *Phytochemistry* **16**: 1503-1505.
- Lee, J., Reeves, R. D., Brooks, R. R. and Jaffré, T. (1978). The relation between nickel and citrate in some nickel accumulating plants. *Phytochemistry* **17**: 1033-1035.
- Lindsay, W. L. and Norvell, W. A. (1978). Development of a DTPA soil test for zinc, iron, manganese and copper. *Journal of the Soil Science Society of America* **42**, 421-428.
- Llugany, M. E., Lombini, A., Poschenrieder, C., Dinelli, E. and Barcello, J. (2003). Different mechanisms account for enhanced copper resistance in *Silene armeria* ecotypes from mine spoil and serpentinite sites. *Plant and Soil* **251**: 55-63.
- Macnair, M. R. and Baker, A. J. M. (1994). Metal-tolerant plants: An evolutionary perspective. In: *Plants and the Chemical Elements* (Ed. M. E. Farago), pp 68-85. VCH, Weinheim, Germany.
- Macnair, M. R. and Gardner, M. (1998). The evolution of edaphic endemics. In: *Endless Forms: Species and Speciation* (Eds D. J. Howard & S. H. Berlocher), pp. 157-171. Oxford University Press, New York, USA.
- Madhok, O. P. and Walker, R. B. (1969). Magnesium nutrition of two species of sunflower. *Plant Physiology* **44**: 1016-1022.
- Mason, H. (1946). The edaphic factor in narrow endemism. The nature of environmental influences. *Madroño* **8**: 209-226.
- Mathys, W. (1977). The role of malate, oxalate, and mustard oil glucosides in the evolution of zinc-resistance in herbage plants. *Physiologia Plantarum* **40**: 130-136.
- Mehta, A. and Farago, M. E. (1994). Metal ions and plant nutrition. In: *Plants and the Chemical Elements* (Ed. M. E. Farago), pp. 32-66. VCH, Weinheim, Germany.
- Mesjasz-Przybyłowicz, J. and Przybyłowicz, W. J. (2001). Phytophagous insects associated with the Ni-hyperaccumulating plant *Berkheya coddii* (Asteraceae) in Mpumalanga, South Africa. *South African Journal of Science* **97**: 596-598.
- Munasinghe, T. and Dissanayake, C. B. (1979). Is the highland -eastern Vijayan boundary in Sri Lanka a possible mineralized belt? *Economic Geology and the Bulletin of the Society of Economic Geologists* **74**: 1495-1496.
- Munasinghe, T. and Dissanayake, C. B. (1980). Is the highland -eastern Vijayan boundary in Sri Lanka a possible mineralized belt ? - a reply. ?. *Economic Geology and the Bulletin of the Society of Economic Geologists* **75**: 775-777.
- Myers, N. (1990). The biodiversity challenge: expanded hot-spots analysis. *The Environmentalist* **10**: 243-256.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B. and Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature* **403**: 853-858.
- Nagy, L. and Proctor, J. (1997). Soil Mg and Ni as causal factors of plant occurrence and distribution at the Meikle Kilrannoch ultramafic site in Scotland. *New Phytologist* **135**: 561-566.
- Novak, F. A. (1928). Quelques remarques relatives au problème de la végétation sur les terrains serpentiniques. *Preslia* **6**: 42-71.
- Pernas, M., Sanchez-Monge, R. and Salcedo, G. (2000). Biotic and abiotic stress can induce cystatin expression in chestnut. *Federation of European Biochemical Societies Letters* **467**: 206-210.
- Pollard, A. J. (2000). Metal hyperaccumulation: a model system for coevolutionary studies. *New Phytologist* **146**: 179-181.
- Pollard, A. J., Powell, K. D., Harper, F. A. and Smith, J. A. C. (2002). The genetic basis for metal hyperaccumulation in plants. *Critical Reviews in Plant Sciences* **21**: 539-566.
- Proctor, J. and Whitten, K. (1971). A population

- of the Valley Pocket Gopher (*Thomomys bottae*) on a serpentine soil. *American Midland Naturalist* **85**: 517-521.
- Proctor, J. and Woodell, S. R. J. (1975). The ecology of serpentine soils. *Advances in Ecological Research* **9**: 255-366.
- Proctor, J. and McGowan, I. D. (1976). Influence of magnesium on nickel toxicity. *Nature* **134**: 260.
- Proctor, J. (2003). Vegetation and soil and plant chemistry on ultramafic rocks in the tropical Far East. *Perspectives in Plant Ecology, Evolution, and Systematics* **6**: 105-124.
- Rajakaruna, N. and Bohm, B. A. (1999). The edaphic factor and patterns of variation in *Lasthenia californica* (Asteraceae). *American Journal of Botany* **86**: 1576-1596.
- Rajakaruna, N. and Bohm, B. A. (2002). Serpentine and its vegetation: a preliminary study from Sri Lanka. *Journal of Applied Botany* **76**: 20-28
- Rajakaruna, N., Harris, C. S. and Towers, G. H. N. (2002b). Antimicrobial activity of plants collected from serpentine outcrops in Sri Lanka. *Pharmaceutical Biology* **40**: 235-244
- Rajakaruna, N. and Whitton, J. (2004). Trends in the evolution of edaphic specialists with an example of parallel evolution in the *Lasthenia californica* complex. In: *Plant adaptation: Molecular Biology and Ecology* (Eds. Q.C.B. Cronk, I.E.P. Taylor, R. Ree and J. Whitton). In press.
- Ranasinghe, N. S. (1987). Serpentinities associated with the precambrian of Sri Lanka. Geological Society of Sri Lanka Special Publication No. 3. Geological Survey Department, Colombo, Sri Lanka.
- Rausser, W. E. (1995). Phytochelatins and related peptides. Structure, biosynthesis, and function. *Plant Physiology* **109**: 1141-1149.
- Reeves, R. D. (1992). The hyperaccumulation of nickel by serpentine plants. In: *The Vegetation of Ultramafic (Serpentine) Soils*. Proceedings of the First International Conference on Serpentine Ecology (Eds. A. J. M. Baker, J. Proctor and R. D. Reeves), pp. 253-277. Intercept Ltd., Hampshire, UK.
- Reeves, R. D. and Baker, A. J. M. (2000). Metal-accumulating plants. In: *Phytoremediation of Toxic Metals: Using Plants to Clean Up the Environment* (Ed. I. Raskin and B. D. Ensley), pp. 193-229. John Wiley and Sons Inc., NY, USA.
- Reeves, R. D., Baker, A. J. M., Borhidi, A. and Berazain, R. (1999). Nickel hyperaccumulation in serpentine flora of Cuba. *Annals of Botany* **83**: 29-38.
- Reeves, R. D. (2003). Tropical hyperaccumulators of metals and their potential for phytoextraction. *Plant and Soil* **249**: 57-65.
- Roberts, B. A. and Proctor, J. (1992). *The Ecology of Areas with Serpentinized Rocks. A World View*. Kluwer Academic Publications, Dordrecht, The Netherlands. 427 p.
- Robertson, A. I. (1992). The relation of nickel toxicity to certain physiological aspects of serpentine ecology: some facts and a new hypothesis. In: *The Vegetation of Ultramafic (Serpentine) Soils*. Proceedings of the First International Conference on Serpentine Ecology (Eds. A. J. M. Baker, J. Proctor and R. D. Reeves), pp 331-336. Intercept Ltd., Hampshire, UK.
- Robinson, N. J. (1990). Metal-binding polypeptides in plants. In: *Heavy Metal Tolerance in Plants: Evolutionary Aspects* (Ed. A. J. Shaw), pp 195-214. CRC Press, Boca Raton, FL, USA..
- Robinson, N. J., Tommey, A. M., Kuske, C. and Jackson, P. J. (1993). Plant metallothioneins. *Journal of Biochemistry* **295**: 1-10.
- Roy, S. (1974). Geobotany in the exploration of Ni in the ultramafics of the Sukinda Valley Orissa. *The Quarterly Journal of the Geological, Mining, and Metallurgical Society of India* **46**: 251-256.

- Salt, D. E., Thurman, D. A., Tomsett, A. B. and Sewell, A. (1989). Copper phytochelatins of *Mimulus guttatus*. *Proceedings of the Royal Society of London, B* **236**: 79.
- Schemske, D. W., Bierzychudek, P. (2001). Perspective: Evolution of flower color in the desert annual *Linanthus parryae*: Wright revisited. *Evolution* **55**: 1269-1282.
- Shaw, A. J. (Ed.) (1990). *Heavy Metal Tolerance in Plants: Evolutionary Aspects*. CRC Press, Boca Raton, FL, USA., 355 p.
- Shaw, A. J. (1999). The evolution of heavy metal tolerance in plants: adaptations, limits, and costs. In: *Genetics and Ecotoxicology. Current Topics in Ecotoxicology and Environmental Chemistry* (Ed. V. E. Forbes), pp 9-30. Taylor and Francis, Inc., Philadelphia, PA, USA.
- Strange, J. and Macnair, M. R. (1991). Evidence for a role for the cell membrane in copper tolerance of *Mimulus guttatus* Fischer Ex DC. *New Phytologist* **119**: 383-388.
- Tadros, T. M. (1957). Evidence for the presence of an edapho-biotic factor in the problem of serpentine tolerance. *Ecology* **38**: 14-23.
- Tilstone, G. H., Macnair, M. R. and Smith, S. E. (1997). Does copper tolerance give cadmium tolerance in *Mimulus guttatus*. *Heredity* **79**, 445-452.
- Tilstone, G. H. and Macnair, M. R. (1997a). Nickel tolerance and copper-nickel co-tolerance in *Mimulus guttatus* from copper mine and serpentine habitats. *Plant and Soil* **191**: 173-180.
- Tilstone, G. H. and Macnair, M. R. (1997b). The consequence of selection for copper tolerance on the uptake and accumulation of copper in *Mimulus guttatus*. *Annals of Botany* **80**: 747-751.
- Verkleij, J. A. C. and Schat, H. (1990). Mechanisms of metal tolerance in higher plants. In: *Heavy Metal Tolerance in Plants: Evolutionary Aspects* (Ed. A. J. Shaw), pp. 179-193. CRC Press, Boca Raton, FL, USA.
- Vlamis, J. and Jenny, H. (1948). Calcium deficiency in serpentine soils as revealed by absorbent technique. *Science* **107**, 549-551.
- Wall, M. A. and Boyd, R. S. (2002). Nickel accumulation in serpentine arthropods from the Red Hills, California. *Pan-Pacific Entomologist* **78**: 168-176.
- Walker, R. B. (1954). The ecology of serpentine soils. II. Factors affecting plant growth on serpentine soils. *Ecology* **35**: 259-266
- Walker, R. B., Walker, H. M., and Ashworth, P. R. (1955). Calcium-magnesium nutrition with special reference to serpentine soils. *Plant Physiology* **30**: 214-221.
- Whiting, S. N., Reeves, R. D. and Baker, A. J. M. (2002). Conserving biodiversity: mining, metallophytes and land reclamation: conservation of biodiversity. *Mining Environmental Management* **10**: 11-16.
- Whiting, S. N., Neumann, P. M., and Baker, A. J. M. (2003). Nickel and zinc Hyperaccumulation by *Alyssum murale* and *Thlaspi caerulescens* (Brassicaceae) do not enhance survival and whole-plant growth under drought stress. *Plant, Cell and Environment* **26**: 351-360.
- Whiting, S. N., Reeves, R. D., Richards, D., Johnson, M. S., Cooke, J. A., Malaisse, F., Paton, A., Smith, J. A. C., Angle, J. S., Chaney, R. L., Ginocchio, R., Jaffré, T., Johns, R., McIntyre, T., Purvis, O. W., Salt, D. E., Schat, H., Zhao, F. J. and Baker, A. J. M. (2004). Research priorities for conservation of metallophytes and their potential for restoration and site remediation. *Restoration Ecology* (in press).
- Whittaker, R. H., Walker, R. B. and Kruckeberg, A. R. (1954). The ecology of serpentine soils: A symposium, Parts I-IV. *Ecology* **35**: 258-288.
- Wild, H. and Bradshaw, A. D. (1977). The evolutionary effects of metalliferous and other anomalous soils in south central Africa. *Evolution* **31**, 282-293.

Zenk, M. H. (1996). Heavy metal detoxification in higher plants - a review. *Gene* **179**: 21-30.

Zhou, J. and Goldsbrough, P. B. (1994) Functional homologs of fungal metallothionein genes from *Arabidopsis*. *Plant Cell* **6**: 875-884.